

Research Article

Geometric Sensitivity of a Pinhole Collimator

Howard Jacobowitz¹ and Scott D. Metzler²

¹ Department of Mathematical Sciences, Rutgers University, Camden, NJ 08102, USA

² Department of Radiology, University of Pennsylvania, Philadelphia, PA 19104, USA

Correspondence should be addressed to Howard Jacobowitz, jacobowi@camden.rutgers.edu

Received 25 August 2009; Revised 7 December 2009; Accepted 19 February 2010

Academic Editor: Harvinder S. Sidhu

Copyright © 2010 H. Jacobowitz and S. D. Metzler. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Geometric sensitivity for single photon emission computerized tomography (SPECT) is given by a double integral over the detection plane. It would be useful to be able to explicitly evaluate this quantity. This paper shows that the inner integral can be evaluated in the situation where there is no gamma ray penetration of the material surrounding the pinhole aperture. This is done by converting the integral to an integral in the complex plane and using Cauchy's theorem to replace it by one which can be evaluated in terms of elliptic functions.

1. Introduction

Nuclear-medicine imaging provides images that assess how the body is functioning [1, 2], as opposed to anatomical modalities (e.g., X-ray computed tomography, commonly known as a CT or "CAT" scans) that provide little or no information about function, but great detail of the body's structure. Nuclear-medicine images the biodistribution of radiolabeled molecules that are typically intravenously injected into the patient in tracer (i.e., nonpharmacological) quantities. The compounds may have different biochemical properties that affect the biodistribution and, hence, the choice of pharmaceuticals used to assess the disease state of a patient.

Two common nuclear-medicine techniques are single photon emission computed tomography (SPECT) and positron emission tomography (PET). Molecules labeled with a SPECT tracer emit a single photon. PET tracers emit a positron, which annihilates with a nearby electron to produce two nearly back-to-back photons at 511 keV, the rest energy of an electron. The line of these two photons contains the emission point of the positron. Many photon pairs are detected in coincidence and reconstructed into a three-dimensional (3D) image of the tracer's distribution.

Since SPECT tracers emit only one photon, the detection of the photon itself gives little information about the location of the emission since neither the direction nor origin




Figure 1: Perspective View of the Knife-edge Pinhole Collimator. A point source at $z = h$ and angle θ emits a photon that intersects the $z = 0$ plane at $(\rho \cos \beta, \rho \sin \beta, 0)$ with an incident angle θ_a . The acceptance angle of the collimator is α and the diameter of the aperture is d . ΔL is the path length of the photon in the attenuating medium. Reproduced from [3] with permission (© 2001 IEEE).

of the emission is known. Consequently, SPECT uses collimation that allows only photons along certain lines to be detected. The origin is still unknown, but the 3D image may be reconstructed when photons from numerous angles are detected [4, 5]. Pinhole collimation is often used when a small organ (e.g., the thyroid) is imaged by a large detector with a SPECT tracer. Figure 1 is a diagram of the apparatus. It is not hard to see that the pinhole collimation's geometric efficiency, namely, the fraction of emitted photons that pass through the circular aperture of the collimator, is approximately given by

$$g = \frac{d^2 \sin^3 \theta}{16h^2}, \quad (1.1)$$

where d is the diameter of the pinhole, h is the perpendicular distance, and θ is the incidence angle of the photon on the aperture plane at the center of aperture. This formula becomes exact in the limit that d goes to zero. Also, an exact formula is easily obtained when $\theta = \pi/2$

$$g\left(\theta = \frac{\pi}{2}\right) = \frac{1}{2} - \frac{1}{2} \left[1 + \left(\frac{d}{2h} \right)^2 \right]^{-1/2} \approx \frac{1}{2} - \frac{1}{2} \left[1 - \frac{1}{2} \left(\frac{d}{2h} \right)^2 \right] = \frac{d^2}{16h^2}. \quad (1.2)$$

We derive stronger versions of these two results in the next section. But the main goal of this paper is to simplify the integral representation for the geometric sensitivity without taking

d or $\theta - \pi/2$ to be small. We are concerned with the case where there is no penetration in the sense that not all gamma rays are stopped by the attenuating material surrounding the pinhole aperture. We take $\theta \neq \pi/2$. The geometric sensitivity, as defined above, is given by the integral over the aperture of the flux times the sine of the incidence angle at that point on the aperture:

$$g = \frac{1}{4\pi h^2} \int_0^{d/2} \rho d\rho \int_0^{2\pi} d\beta \sin^3 \theta_a, \quad (1.3)$$

where ρ and β are polar coordinates on the aperture plane and θ_a is the incidence angle of the photon at a particular value of ρ and β , and is given by [3]

$$\cot^2 \theta_a = \cot^2 \theta - 2 \frac{\rho}{h} \cot \theta \cos(\beta - \phi) + \frac{\rho^2}{h^2}, \quad (1.4)$$

where ϕ is the third coordinate of the point source. For integrals over 2π the difference $\beta - \phi$ can be replaced with only β without loss of generality. This yields

$$\sin^2 \theta_a = \left[\csc^2 \theta - 2 \frac{\rho}{h} \cot \theta \cos \beta + \frac{\rho^2}{h^2} \right]^{-1}. \quad (1.5)$$

Applying this to (1.3),

$$g = \frac{1}{4\pi h^2} \int_0^{d/2} \rho d\rho \int_0^{2\pi} d\beta \left[\csc^2 \theta - 2 \frac{\rho}{h} \cot \theta \cos \beta + \frac{\rho^2}{h^2} \right]^{-3/2}. \quad (1.6)$$

When there is penetration the above would be modified to

$$g = \frac{1}{4\pi h^2} \int_0^\infty \rho d\rho \int_0^{2\pi} d\beta \left[\csc^2 \theta - 2 \frac{\rho}{h} \cot \theta \cos \beta + \frac{\rho^2}{h^2} \right]^{-3/2} e^{-\mu \Delta L}, \quad (1.7)$$

where the path length through attenuating material, ΔL , is zero for $\rho < d/2$ and is otherwise given by (11) in [3].

Thus, (1.7) can be rewritten as

$$\begin{aligned} g &= g_{\text{NP}} + \frac{1}{4\pi h^2} \int_{d/2}^\infty \rho d\rho \int_0^{2\pi} d\beta \left[\csc^2 \theta - 2 \frac{\rho}{h} \cot \theta \cos \beta + \frac{\rho^2}{h^2} \right]^{-3/2} e^{-\mu \Delta L} \\ &= g_{\text{NP}} + g_{\text{PEN}}, \end{aligned} \quad (1.8)$$

where g_{PEN} is the efficiency due to penetration and g_{NP} , the efficiency under the assumption of no penetration, is given by (1.6).

In this paper we use complex variable methods to calculate the β integral in (1.6).

2. Two Approximations for g_{NP}

The special form of g_{NP} makes it easy to derive approximations for d/h small and for $\theta - \pi/2$ small which generalize (1.1) and (1.2). Before doing this, we observe that only even powers of d and of $\theta - \pi/2$ can occur in the expansion of g_{NP} . We see this by writing the integral as the sum of integrals from 0 to π and from π to 2π .

2.1. The Expansion for d Small

Let us, in this section, denote g_{NP} by g . Upon substituting $r = d/2h$ and $\eta = \rho/h$ into (1.6) we obtain

$$\begin{aligned} g(r, \theta) &= \frac{1}{4\pi} \int_0^{2\pi} \int_0^r \left[\csc^2 \theta - 2\eta \cot \theta \cos \beta + \eta^2 \right]^{-3/2} \eta \, d\eta \, d\beta, \\ &= \frac{1}{4\pi} \int_0^{2\pi} \left(\int_0^r \eta \phi(\beta, \eta, \theta) \, d\eta \right) d\beta. \end{aligned} \quad (2.1)$$

Clearly, $g(0, \theta) = 0$. Since

$$g_r(r, \theta) = \frac{1}{4\pi} \int_0^{2\pi} r \phi(\beta, r, \theta) \, d\beta, \quad (2.2)$$

we have $g_r(0, \theta) = 0$. Differentiating (2.12) three more times with respect to r we obtain

$$\begin{aligned} g_{rr}(r, \theta) &= \frac{1}{4\pi} \int_0^{2\pi} (r \phi_r(\beta, 0, \theta) + \phi(\beta, r, \theta)) \, d\beta, \\ g_{rrr}(r, \theta) &= \frac{1}{4\pi} \int_0^{2\pi} (r \phi_{rr}(\beta, 0, \theta) + 2\phi_r(\beta, r, \theta)) \, d\beta, \\ g_{rrrr}(r, \theta) &= \frac{1}{4\pi} \int_0^{2\pi} (r \phi_{rrr}(\beta, 0, \theta) + 3\phi_{rr}(\beta, r, \theta)) \, d\beta, \end{aligned} \quad (2.3)$$

and then setting $r = 0$ in each of the resulting equations we obtain

$$\begin{aligned} g_{rr}(0, \theta) &= \frac{1}{4\pi} \int_0^{2\pi} \phi(\beta, 0, \theta) \, d\beta, \\ g_{rrr}(0, \theta) &= \frac{1}{2\pi} \int_0^{2\pi} \phi_r(\beta, 0, \theta) \, d\beta, \\ g_{rrrr}(0, \theta) &= \frac{3}{4\pi} \int_0^{2\pi} \phi_{rr}(\beta, 0, \theta) \, d\beta. \end{aligned} \quad (2.4)$$

For convenience let us write

$$u = \csc^2\theta, \quad v = \cot\theta, \quad (2.5)$$

so ϕ becomes

$$\phi(\beta, r, \theta) = \left(u - 2vr \cos \beta + r^2\right)^{-3/2}. \quad (2.6)$$

Then

$$\begin{aligned} \phi(\beta, 0, \theta) &= u^{-3/2}, \\ \phi_r(\beta, 0, \theta) &= 3u^{-5/2}v \cos \beta, \\ \phi_{rr}(\beta, 0, \theta) &= 15u^{-7/2}v^2 \cos^2 \beta - 3u^{-5/2}. \end{aligned} \quad (2.7)$$

We substitute into the Taylor expansion

$$g(r, \theta) = g(0, \theta) + g_r(0, \theta)r + \frac{1}{2!}g_{rr}(0, \theta)r^2 + \frac{1}{3!}g_{rrr}(0, \theta)r^3 + \frac{1}{4!}g_{rrrr}(0, \theta)r^4 + \dots \quad (2.8)$$

and obtain

$$\begin{aligned} g(r, \theta) &= \frac{1}{2} \left(\frac{1}{4\pi} \int_0^{2\pi} u^{-3/2} d\beta \right) r^2 + \frac{1}{3!} \left(\frac{3}{2\pi} \int_0^{2\pi} 6u^{-5/2}v \cos \beta d\beta \right) r^3 \\ &+ \frac{1}{4!} \left(\frac{1}{4\pi} \int_0^{2\pi} 3(15u^{-7/2}v^2 \cos^2 \beta - 3u^{-5/2}) d\beta \right) r^4 + \dots \end{aligned} \quad (2.9)$$

Since

$$\int_0^{2\pi} \cos \beta d\beta = 0, \quad \int_0^{2\pi} \cos^2 \beta d\beta = \pi \quad (2.10)$$

and $u^{-1} = \sin^2\theta$ we obtain

$$g(r, \theta) = \frac{r^2}{4} \sin^3\theta + \frac{3}{32} (5\sin^5\theta \cos^2\theta - 2\sin^5\theta) r^4 + \dots \quad (2.11)$$

Finally we replace $\cos^2\theta$ by $1 - \sin^2\theta$ and r by $d/2h$ and obtain

$$g_{\text{NP}} = \frac{1}{16} \sin^3\theta \left(\frac{d}{h}\right)^2 + \frac{3}{512} \sin^5\theta (3 - 5\sin^2\theta) \left(\frac{d}{h}\right)^4 + \mathcal{O}(d^6). \quad (2.12)$$

2.2. The Expansion for $\theta - \pi/2$ Small

To start we let

$$f(\theta, \eta, \beta) = \left(\csc^2 \theta - 2\eta \cot \theta \cos \beta + \eta^2 \right)^{-3/2} \quad (2.13)$$

and compute its Taylor expansion at $\theta = \pi/2$,

$$\begin{aligned} f(\theta, \eta, \beta) &= \left(1 + \eta^2\right)^{-3/2} - 3\left(1 + \eta^2\right)^{-5/2} \eta \cos \beta \left(\theta - \frac{\pi}{2}\right) \\ &+ \frac{1}{2!} \left(15\left(1 + \eta^2\right)^{-7/2} \eta^2 \cos^2 \beta - 3\left(1 + \eta^2\right)^{-5/2}\right) \left(\theta - \frac{\pi}{2}\right)^2 \\ &+ \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^3. \end{aligned} \quad (2.14)$$

Since

$$g_{\text{NP}} = \frac{1}{4\pi} \int_0^{2\pi} \int_0^{d/2h} f(\theta, \eta, \beta) \eta d\eta d\beta \quad (2.15)$$

and since we know that only even powers of $\theta - \pi/2$ occur, we obtain

$$g_{\text{NP}} = \frac{1}{2} \left(1 - \frac{1}{\sqrt{1 + (d/2h)^2}}\right) - \frac{3}{8} \frac{(d/2h)^2}{\left(1 + (d/2h)^2\right)^{5/2}} \left(\theta - \frac{\pi}{2}\right)^2 + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4. \quad (2.16)$$

3. The Contour Integral

We have

$$I = \int_0^{2\pi} d\beta \left[\csc^2 \theta - 2\frac{\rho}{h} \cot \theta \cos \beta + \frac{\rho^2}{h^2} \right]^{-3/2}, \quad (3.1)$$

$$g_{\text{NP}} = \frac{1}{4\pi h^2} \int_0^{d/2} I \rho d\rho. \quad (3.2)$$

I may be recast as a contour integral by setting $z = e^{i\beta}$:

$$\begin{aligned} I &= \int_{\Gamma} \frac{dz}{iz} \left[\frac{-b}{2z} \left(z^2 - 2\frac{a}{b}z + 1 \right) \right]^{-3/2} \\ &= \left(\frac{2}{b}\right)^{3/2} \frac{1}{i} \int_{\Gamma} \left[\frac{(z - x_0)(x_1 - z)}{z} \right]^{-3/2} \frac{dz}{z}, \end{aligned} \quad (3.3)$$

where

$$\begin{aligned}
 &\Gamma \text{ is the unit circle,} \\
 &a = \csc^2 \theta + \frac{\rho^2}{h^2}, \\
 &b = 2 \frac{\rho}{h} \cot \theta, \\
 &x_0 = \frac{a}{b} \left[1 - \sqrt{1 - \frac{b^2}{a^2}} \right], \\
 &x_1 = \frac{a}{b} \left[1 + \sqrt{1 - \frac{b^2}{a^2}} \right].
 \end{aligned} \tag{3.4}$$

We note that $a > b$ and so $0 < x_0 < 1 < x_1$.

In this section we take ρ, θ , and h to be constant. We write

$$\begin{aligned}
 F(z) &= \left[\frac{z}{(z-x_0)(x_1-z)} \right] \left[\frac{z}{((z-x_0)(x_1-z))} \right]^{1/2} \left[\frac{1}{z} \right] \\
 &= \frac{1}{(z-x_0)(x_1-z)} \left[\frac{z}{(z-x_0)(x_1-z)} \right]^{1/2}, \\
 I &= \left(\frac{2}{b} \right)^{3/2} \frac{1}{i} \int_{\Gamma} F(z) dz.
 \end{aligned} \tag{3.5}$$

We need to be careful because of the square roots. We use the usual convention that

$$\lim_{y \rightarrow 0^\pm} \sqrt{-1 + iy} = \pm i. \tag{3.6}$$

We wish to show that $F(z)$ is holomorphic (and single valued) in

$$\{z : |z| < 1\} - \{x : 0 < x < x_0\}. \tag{3.7}$$

To isolate the square root factor in F we set

$$g(z) = \frac{z}{(z-x_0)(x_1-z)}. \tag{3.8}$$

We take $|z| \leq 1$. If in addition $\operatorname{Re} z < 0$ and $|\operatorname{Im} z| \ll |z - x_0|$ then $\operatorname{Re} (z - x_0)(x_1 - z) < 0$. This implies that $g(z)$ never takes on negative real values and so has a single valued square root in $\operatorname{Re} z < 0$. Similarly, if $|z| \leq 1$, and $\operatorname{Re} z > x_0$ (no condition on $\operatorname{Im} z$ now) then $\operatorname{Re} (z - x_0)(x_1 - z) > 0$ and $g(z)$ has a single valued square root in $\operatorname{Re} z > x_0$.




Figure 2: The curve γ .

It follows from Cauchy's theorem that the integral of $F(z)$ over the unit circle Γ is equal to the integral over the "barbell" γ around 0 and x_0 . More explicitly, γ is the boundary of

$$B = B_\epsilon(0) \cup B_\epsilon(x_0) \cup R, \quad (3.9)$$

where $B_\epsilon(x)$ is the ball of radius ϵ and centered at x and R is the rectangle

$$R = \{z = x + iy : 0 \leq x \leq x_0, -\delta \leq y \leq \delta\} \quad (3.10)$$

with $\delta \ll \epsilon$, see Figure 2.

The curve γ decomposes into smooth pieces which we describe using the angle $\alpha = \arcsin(\delta/\epsilon)$

$$\begin{aligned} \gamma_r &= \{z : |z - x_0| = \epsilon, -\pi + \alpha \leq \arg z - x_0 \leq \pi - \alpha\}, \\ \gamma_l &= \{z : |z| = \epsilon, \alpha \leq \arg(z) \leq 2\pi - \alpha\}, \\ \gamma_\pm &= \{z : z = x \pm i\delta, \epsilon \cos \alpha \leq x \leq x_0 - \epsilon \cos \alpha\}. \end{aligned} \quad (3.11)$$

To bound the integral over γ_l we set $z = \epsilon e^{i\theta}$

$$\begin{aligned} \left| \int_{\gamma_l} F(z) dz \right| &= \left| \int_{\alpha}^{2\pi - \alpha} F(\epsilon e^{i\theta}) \epsilon e^{i\theta} d\theta \right| \\ &= \mathcal{O}(\epsilon^{3/2}). \end{aligned} \quad (3.12)$$

To evaluate $\int_{\gamma_r} F(z) dz$ we set $z = x_0 + \epsilon e^{i\theta}$. Then

$$F(x_0 + \epsilon e^{i\theta}) = C \epsilon^{-3/2} e^{-(3/2)i\theta} + \mathcal{O}(\epsilon^{-1/2}) \quad (3.13)$$

with

$$C = \frac{x_0^{1/2}}{(x_1 - x_0)^{3/2}}, \quad (3.14)$$

and so

$$\begin{aligned} \int_{\gamma_r} F(z)dz &= \int_{-\pi+\alpha}^{\pi-\alpha} \left(\epsilon^{-3/2} C e^{-(3/2)i\theta} + \mathcal{O}(\epsilon^{-1/2}) \right) \epsilon i e^{i\theta} (1 + \mathcal{O}(\epsilon)) d\theta \\ &= \int_{-\pi+\alpha}^{\pi-\alpha} C i \epsilon^{-1/2} e^{-(1/2)i\theta} + \mathcal{O}(\epsilon^{1/2}) d\theta. \end{aligned} \tag{3.15}$$

Since $\alpha = \mathcal{O}(\epsilon)$, we may set $\alpha = 0$ and absorb the error in the $\mathcal{O}(\epsilon^{1/2})$ term. We then do the integration and obtain

$$\int_{\gamma_r} F(z)dz = \frac{4iC}{\sqrt{\epsilon}} + \mathcal{O}(\epsilon^{1/2}). \tag{3.16}$$

We want to compute the integral of F over γ_+ . We need to be careful about the square root. Recall that we are assuming

$$\epsilon \cos \alpha < x < x_0 - \epsilon \cos \alpha, \quad \delta \ll \epsilon. \tag{3.17}$$

Lemma 3.1. *On γ_+*

$$F(z) = i((x_0 - x)(x_1 - x))^{-3/2} x^{1/2} + \mathcal{O}(\delta) \tag{3.18}$$

and on γ_-

$$F(z) = -i((x_0 - x)(x_1 - x))^{-3/2} x^{1/2} + \mathcal{O}(\delta). \tag{3.19}$$

Proof. This is a consequence of the branching of the square root function. Let

$$\begin{aligned} H(z) &= \frac{z}{(z - x_0)(x_1 - z)} = \frac{z(\bar{z} - x_0)(x_1 - \bar{z})}{|z - x_0|^2 |x_1 - z|^2}, \\ h(z) &= z(\bar{z} - x_0)(x_1 - \bar{z}). \end{aligned} \tag{3.20}$$

We see that

$$h(x + i\delta) = x(x - x_0)(x_1 - x) + i\delta(x^2 - 1) + \mathcal{O}(\delta^2). \tag{3.21}$$

Recall that δ is replaced by $-\delta$ on γ_- and that $0 < x < x_0 < x_1$. So $\text{Re } H(z) < 0$ on both γ_+ and γ_- while $\text{Im } H(z) < 0$ on γ_+ and $\text{Im } H(z) > 0$ on γ_- . Thus on γ_+

$$\begin{aligned} H(z)^{1/2} &= -i \left| \frac{x}{(x - x_0)(x_1 - x)} \right|^{1/2} + \mathcal{O}(\delta) \\ &= -i \left(\frac{x}{(x_0 - x)(x_1 - x)} \right)^{1/2} + \mathcal{O}(\delta), \end{aligned} \tag{3.22}$$

while on γ_-

$$H(z)^{1/2} = i \left(\frac{x}{(x_0 - x)(x_1 - x)} \right)^{1/2} + \mathcal{O}(\delta). \quad (3.23)$$

Since

$$\begin{aligned} F(z) &= F(x) + \mathcal{O}(\delta) = \frac{1}{(x - x_0)(x_1 - x)} H(z)^{1/2} + \mathcal{O}(\delta) \\ &= -\frac{1}{(x_0 - x)(x_1 - x)} H(z)^{1/2} + \mathcal{O}(\delta) \\ &= -\frac{1}{(x_0 - x)(x_1 - x)} \left(-i \left(\frac{x}{(x_0 - x)(x_1 - x)} \right)^{1/2} \right) + \mathcal{O}(\delta), \end{aligned} \quad (3.24)$$

the lemma follows. \square

We also need to be careful about the orientation. The curve γ is oriented counterclockwise. This means that the integration over γ_+ goes in the direction of x_0 to 0. We need to reverse its sign in order to write it in the conventional way

$$\begin{aligned} \int_{\gamma_+} F(z) dz &= - \int_{\epsilon \cos \alpha}^{x_0 - \epsilon \cos \alpha} F(x + i\delta) dx \\ &= - \int_{\epsilon \cos \alpha}^{x_0 - \epsilon \cos \alpha} \frac{ix^{1/2} dx}{(x_0 - x)^{3/2} (x_1 - x)^{3/2}} + \mathcal{O}(\epsilon). \end{aligned} \quad (3.25)$$

We let $x = x_0 t$ and $\Delta = \epsilon x_0^{-1} \cos \alpha$ and use $\delta \ll \epsilon$ to obtain

$$\int_{\gamma_+} F(z) dz = - \int_{\Delta}^{1-\Delta} \frac{ix_0^{3/2} t^{1/2} dx}{(x_0 - x_0 t)^{3/2} (x_1 - x_0 t)^{3/2}} + \mathcal{O}(\epsilon). \quad (3.26)$$

Now let

$$\tau = \frac{x_1}{x_0} > 1. \quad (3.27)$$

We have

$$\int_{\gamma_+} F(z) dz = -\frac{i}{x_0^{3/2}} \int_{\Delta}^{1-\Delta} \frac{t^{1/2} dt}{(\tau - t)^{3/2} (1 - t)^{3/2}} + \mathcal{O}(\epsilon). \quad (3.28)$$

We may let $\Delta \rightarrow 0$ in the lower limit, but not in the upper limit. So we now have

$$\int_{\gamma_+} F(z)dz = -\frac{i}{x_0^{3/2}} \int_0^{1-\Delta} \frac{t^{1/2} dt}{(\tau-t)^{3/2}(1-t)^{3/2}} + \mathcal{O}(\epsilon). \tag{3.29}$$

Here $\Delta = \epsilon x_0^{-1} \cos \alpha$. We may let $\delta \rightarrow 0$ and redefine Δ to be ϵx_0^{-1} .

We get the same value for the integral over γ_- . This is because the integrand is the negative of that in (3.29) but the orientation is reversed and so the two negatives cancel.

For any function $g(t)$, continuously differentiable on the interval $[0, 1]$, we have

$$\int_0^{1-\Delta} \frac{g(t)}{(1-t)^{3/2}} dt = 2\frac{g(1)}{\Delta^{1/2}} - 2g(0) - 2\int_0^1 (1-t)^{-1/2} g'(t) dt + \mathcal{O}(\Delta^{1/2}) \tag{3.30}$$

as can easily be seen by an integration by parts.

We apply this to $g(t) = t^{1/2}/(\tau-t)^{3/2}$ as in (3.29). This yields

$$\begin{aligned} \int_{\gamma_+} F(z)dz &= \frac{-2i}{(\tau-1)^{3/2} x_0^{3/2} \Delta^{1/2}} \\ &+ \frac{i}{x_0^{3/2}} \left(\int_0^1 t^{-1/2} (\tau-t)^{-3/2} + 3t^{1/2} (\tau-t)^{-5/2} dt \right) + \mathcal{O}(\epsilon^{1/2}). \end{aligned} \tag{3.31}$$

We now take twice this quantity and add to it the results of (3.14) and (3.16),

$$\begin{aligned} \int_{|z|=1} F(z)dz &= \int_{\gamma} F(z)dz \\ &= \frac{4iC}{\sqrt{\epsilon}} + 2 \left(\frac{-2i\sqrt{x_0}}{(x_1-x_0)^{-3/2}\sqrt{\epsilon}} \right. \\ &\quad \left. + \frac{i}{x_0^{3/2}} \int_0^1 (1-t)^{-1/2} \left((\tau-t)^{-3/2} t^{-1/2} + 3t^{1/2} (\tau-t)^{-5/2} \right) dt \right) \\ &\quad + \mathcal{O}(\epsilon^{1/2}). \end{aligned} \tag{3.32}$$

We substitute the value for C from (3.14) and let $\epsilon \rightarrow 0$

$$\int_{|z|=1} F(z)dz = \frac{2i}{x_0^{3/2}} \int_0^1 (1-t)^{-1/2} \left((\tau-t)^{-3/2} t^{-1/2} + 3t^{1/2} (\tau-t)^{-5/2} \right) dt. \tag{3.33}$$

Finally, we evaluate this integral using Mathematica and obtain

$$\int_{|z|=1} F(z) dz = \left(\frac{2i}{x_0^{3/2}} \right) \frac{2(2\tau E[1/\tau] - (-1 + \tau)K[1/\tau])}{(\tau - 1)^2 \sqrt{\tau}}, \quad (3.34)$$

where K and E are the complete elliptic functions of the first and second kinds

$$\begin{aligned} K[m] &= \int_0^1 (1 - mt^2)^{-1/2} (1 - t^2)^{-1/2} dt, \\ E[m] &= \int_0^1 (1 - mt^2)^{1/2} (1 - t^2)^{-1/2} dt. \end{aligned} \quad (3.35)$$

We now have

$$\begin{aligned} I &= \left(\frac{2}{b} \right)^{3/2} \frac{1}{i} \int_{\Gamma} F(z) dz \\ &= \left(\frac{2}{b} \right)^{3/2} \frac{2}{x_0^{3/2}} \frac{2(2\tau E(1/\tau) - (\tau - 1)K(1/\tau))}{(\tau - 1)^2 \sqrt{\tau}} \\ &= \left(\frac{2}{b} \right)^{3/2} \frac{4\tau^{1/4}}{(\tau - 1)^2} (2\tau E(1/\tau) - (\tau - 1)K(1/\tau)), \end{aligned} \quad (3.36)$$

$$g_{\text{NP}} = \frac{1}{4\pi h^2} \int_0^{d/2} \rho I d\rho \quad (3.37)$$

with

$$b = 2 \frac{\rho}{h} \cot \theta, \quad a = \csc^2 \theta + \frac{\rho^2}{h^2}, \quad (3.38)$$

$$\tau = \frac{x_1}{x_0} = \frac{1}{x_0^2},$$

$$x_0 = \frac{a}{b} \left[1 - \sqrt{1 - \frac{b^2}{a^2}} \right], \quad (3.39)$$

$$x_1 = \frac{a}{b} \left[1 + \sqrt{1 - \frac{b^2}{a^2}} \right].$$

4. Limiting Cases

We validate (3.36) by considering two limiting cases.

4.1. Expansion for Small d

We look for an expansion of g_{NP} under the assumption that d/h is small and rederive (2.12). The dependence of x_0 on ρ is given in (3.39). The two-term Taylor expansion of x_0 as a function of ρ is

$$x_0 = A\rho + B\rho^3 + \mathcal{O}(\rho^5) \quad (4.1)$$

with

$$A = \frac{\cos\theta \sin\theta}{h}, \quad B = A^3 - \frac{A}{h^2 \csc^2\theta}. \quad (4.2)$$

Since

$$\tau = \frac{1}{x_0^2}, \quad (4.3)$$

we have

$$\frac{1}{\tau} = A^2\rho^2 + \mathcal{O}(\rho^4), \quad (4.4)$$

$$\frac{1}{(\tau-1)^2} = \frac{1}{\tau^2} + \frac{2}{\tau^3} + \mathcal{O}(\rho^8). \quad (4.5)$$

It is convenient to work with the quantity $\rho^2\tau$, so we note, using (4.1),

$$\rho^2\tau = \frac{1}{A^2} - \frac{2B}{A^3}\rho^2 + \mathcal{O}(\rho^4). \quad (4.6)$$

We start with

$$I = \left(\frac{2}{b}\right)^{3/2} \frac{4\tau^{1/4}}{(\tau-1)^2} \left(2\tau E\left(\frac{1}{\tau}\right) - (\tau-1)K\left(\frac{1}{\tau}\right)\right). \quad (4.7)$$

Substituting the expression for b from (3.38) and for $(\tau-1)^2$ from (4.5) we obtain, after some manipulation

$$I = \left(\left(\frac{h}{\cot\theta}\right)^{3/2} \frac{4}{(\rho^2\tau)^{3/4}} \frac{1}{\tau} + \mathcal{O}(\rho^4)\right) \left(2\tau E\left(\frac{1}{\tau}\right) - (\tau-1)K\left(\frac{1}{\tau}\right)\right). \quad (4.8)$$

We now use the expansions (as given by Mathematica)

$$\begin{aligned} E(t) &= \frac{\pi}{2} - \frac{\pi t}{8} + \mathcal{O}(t^2), \\ K(t) &= \frac{\pi}{2} + \frac{\pi t}{8} + \mathcal{O}(t^2). \end{aligned} \quad (4.9)$$

So

$$I = \left(\frac{h}{\cot\theta} \right)^{3/2} \frac{4}{(\rho^2\tau)^{3/4}} \left(\frac{\pi}{2} + \frac{\pi}{8\tau} + \frac{\pi}{\tau} \right) + \mathcal{O}(\rho^4). \quad (4.10)$$

From the expression for $\rho^2\tau$ in (4.6), we derive

$$\frac{1}{(\rho^2\tau)^{3/4}} = A^{3/2} \left(1 + \frac{3}{2} \frac{B}{A} \rho^2 + \mathcal{O}(\rho^4) \right), \quad (4.11)$$

and this leads to

$$I = 4 \left(\frac{h}{\cot\theta} \right)^{3/2} A^{3/2} \left(\frac{\pi}{2} + \frac{\pi}{8\tau} + \frac{\pi}{\tau} + \frac{3B\rho^2\pi}{4A} \right) + \mathcal{O}(\rho^4). \quad (4.12)$$

We now substitute for A and B using (4.2) and also for τ^{-1} using (4.4)

$$I = \sin^3\theta \left(2\pi + \frac{3\pi}{2} \left(3\sin^2\theta - 5\sin^4\theta \right) \frac{\rho^2}{h^2} \right) + \mathcal{O}(\rho^4). \quad (4.13)$$

Substituting this into (3.37) and integrating, we again derive (2.12).

4.2. An Expansion for θ Near $\pi/2$

We show how (4.7) also gives the expansion of Section 2.2. From (3.38) we see that

$$a = 1 + \frac{\rho^2}{h^2} + \left(\theta - \frac{\pi}{2} \right)^2 + \mathcal{O}\left(\theta - \frac{\pi}{2} \right)^4, \quad (4.14)$$

$$b = -\frac{2\rho}{h} \left(\theta - \frac{\pi}{2} \right) + \mathcal{O}\left(\theta - \frac{\pi}{2} \right)^3. \quad (4.15)$$

So from (3.39) we have

$$x_0 = \frac{b}{2a} \left(1 + \frac{b^2}{4a^2} + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^3 \right). \quad (4.16)$$

Thus

$$\tau = \frac{4a^2}{b^2} - 2 + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^2, \quad (4.17)$$

$$\frac{1}{(\tau - 1)^2} = \frac{1}{\tau^2} \left(1 + \frac{2}{\tau} + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4 \right), \quad (4.18)$$

$$\frac{1}{\tau} \left(2\tau E\left(\frac{1}{\tau}\right) - (\tau - 1)K\left(\frac{1}{\tau}\right) \right) = \frac{\pi}{2} + \frac{9\pi}{8\tau} + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4.$$

These expansions let us write (4.7) as

$$\begin{aligned} I &= \left(\frac{2}{b}\right)^{3/2} \frac{4}{\tau^{3/4}} \left(\left(1 + \frac{2}{\tau}\right) \left(\frac{\pi}{2} + \frac{9\pi}{8\tau}\right) + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4 \right) \\ &= \frac{2^{7/2}}{(b^2\tau)^{3/4}} \left(\left(1 + \frac{2}{\tau}\right) \left(\frac{\pi}{2} + \frac{9\pi}{8\tau}\right) + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4 \right). \end{aligned} \quad (4.19)$$

Using (4.15) and (4.17) we obtain a substitute for (4.1)

$$\frac{1}{(b^2\tau)^{3/4}} = 4^{-3/4} a^{-3/2} \left(1 + \frac{3b^2}{8a^2} \right) + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4. \quad (4.20)$$

This gives, in light of (4.14),

$$\begin{aligned} I &= 2\pi a^{-3/2} + \frac{15}{8}\pi b^2 a^{-7/2} + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4 \\ &= 2\pi \left(1 + \frac{\rho^2}{h^2} \right)^{-3/2} - 3\pi \left(1 + \frac{\rho^2}{h^2} \right)^{-5/2} \left(\theta - \frac{\pi}{2} \right)^2 \\ &\quad + \frac{15}{2}\pi \frac{\rho^2}{h^2} \left(1 + \frac{\rho^2}{h^2} \right)^{-7/2} \left(\theta - \frac{\pi}{2} \right)^2 + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4. \end{aligned} \quad (4.21)$$

Finally, we integrate term by term.

$$\begin{aligned}
g &= \frac{1}{4\pi h^2} \int_0^{d/2} \rho I(\rho) d\rho \\
&= \frac{1}{4\pi h^2} \left(2\pi h^2 \left(1 - \left(1 + \left(\frac{d}{2h} \right)^2 \right)^{-1/2} \right) \right. \\
&\quad \left. - 3\pi h^2 \frac{1}{3} \left(1 - \left(1 + \left(\frac{d}{2h} \right)^2 \right)^{-3/2} \right) \left(\theta - \frac{\pi}{2} \right)^2 \right. \\
&\quad \left. + \frac{15}{2} \pi h^2 \left(\frac{2}{15} - \frac{2 + 5(d/2h)^2}{15} \left(1 + \left(\frac{d}{2h} \right)^2 \right)^{-5/2} \right) \left(\theta - \frac{\pi}{2} \right)^2 \right) \quad (4.22) \\
&\quad + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4 \\
&= \frac{1}{2} \left(1 - \left(1 + \left(\frac{d}{2h} \right)^2 \right)^{-1/2} \right) - \frac{3}{8} \left(1 + \left(\frac{d}{2h} \right)^2 \right)^{-5/2} \left(\frac{d}{2h} \right)^2 \left(\theta - \frac{\pi}{2} \right)^2 \\
&\quad + \mathcal{O}\left(\theta - \frac{\pi}{2}\right)^4.
\end{aligned}$$

5. Numerical Confirmation

The result in (3.36) is further validated in this section with numerical integration of (3.1). In this original formulation, the integral depends on the ratio of ρ/h and on the value of θ . We analytically considered one limiting case for each of these quantities in the previous section. We consider these and other values for these quantities in this section using numerical methods. For fixed values of ρ/h , the numerical integration (solid lines) is shown with the results of the residue calculation (open circles) in Figure 3 as a function of θ . Figure 4 shows the numerical integration and residue results for fixed values of θ as a function of ρ/h . All corresponding values of the residue calculation and the numerical integration agree.

6. Conclusions

The accuracy of collimator efficiency is important for design and for quantitative reconstruction programs. Analytic forms obviate the need for numerical models, which do not readily offer insight into the interplay of collimator parameters. This paper approaches the problem by recasting one of the integrals in terms of contour integration. This integration was completed, but the authors were unable to execute the second integral to yield a complete analytic solution. The result was studied for special cases with known outcomes for validation. Future efforts may be able to build on this validated result to find a complete analytic solution.




Figure 3: Numerical integration of (3.1) (solid lines) and residue calculation of (3.36) (open circles) as a function of θ for fixed values of ρ/h : 0.0, 0.2, 0.4, 0.6, 0.8, and 1.0.




Figure 4: Numerical integration of (3.1) (solid lines) and residue calculation of (3.36) (open circles) as a function of ρ/h for fixed values of θ : 50, 60, 70, 80, and 90 degrees.

Acknowledgment

Scott D. Metzler was supported by the National Institute for Biomedical Imaging and Bioengineering of the National Institutes of Health under Grant R01-EB-6558.

References

- [1] M. E. Phelps, "PET: the merging of biology and imaging into molecular imaging," *Journal of Nuclear Medicine*, vol. 41, no. 4, pp. 661–681, 2000.
- [2] S. H. Britz-Cunningham and S. J. Adelstein, "Molecular targeting with radionuclides: state of the science," *Journal of Nuclear Medicine*, vol. 44, no. 12, pp. 1945–1961, 2003.
- [3] S. D. Metzler, J. E. Bowsher, M. F. Smith, and R. J. Jaszczak, "Analytic determination of pinhole collimator sensitivity with penetration," *IEEE Transactions on Medical Imaging*, vol. 20, no. 8, pp. 730–741, 2001.
- [4] S. S. Orlov, "Theory of three dimensional reconstruction. I. Conditions for a complete set of projections," *Soviet Physics Crystallography*, vol. 20, no. 3, pp. 312–314, 1975.
- [5] H. K. Tuy, "An inversion formula for cone-beam reconstruction," *SIAM Journal on Applied Mathematics*, vol. 43, no. 3, pp. 546–552, 1983.